

Math 1301 Practice Final Exam Version 2

Vanderbilt University

8 December 2025

Name: Key

Please do not open the exam until instructed to do so.
You are allowed a one-page (double-sided) formula sheet.
Your instructor may ask to see your formula sheet.
No calculators, phones, computers, smart watches, etc. are permitted.

The Vanderbilt Honor Code applies.

Part 1. (20 points)

Q1 $\frac{d}{dx} [\log_9(5^{\sin(e^x)}) + \cos^{-1}(\ln(\sqrt[50]{x}))] =$

$$\log_9(5) \cos(e^x) e^x - \frac{1}{x \sqrt{1 - 1/50^2 \ln^2(x)}} \cdot \frac{1}{50}$$

$$\begin{aligned} & \log_9(5^{\sin(e^x)}) + \cos^{-1}(\ln(\sqrt[50]{x})) \\ &= \sin(e^x) \log_9(5) + \cos^{-1}\left(\frac{1}{50} \ln(x)\right) \end{aligned}$$

Q2 $\frac{d}{dx} \ln^{5^x}(x) =$

$$\ln^{5^x}(x) \left[5^x \ln(5) \ln(\ln(x)) + \frac{5^x}{x \ln(x)} \right]$$

$$\begin{aligned} y &= \ln^{5^x}(x) \\ \ln(y) &= 5^x \ln(\ln(x)) \\ \frac{1}{y} \frac{dy}{dx} &= 5^x \ln(5) \ln(\ln(x)) + \frac{5^x}{x \ln(x)} \\ \frac{dy}{dx} &= \ln^{5^x}(x) \left[5^x \ln(5) \ln(\ln(x)) + \frac{5^x}{x \ln(x)} \right] \end{aligned}$$

Q3 $\lim_{x \rightarrow 0^+} \left(\frac{\sin(x)}{x} \right)^{1/x^2} =$


$$1/e^{1/6}$$

$$\left(\frac{\sin(x)}{x} \right)^{1/x^2} = e^{\frac{1}{x^2} \ln(\sin(x)) - \frac{1}{x^2} \ln(x)}$$

$$\begin{aligned} \lim_{x \rightarrow 0^+} \frac{\ln(\sin(x)) - \ln(x)}{x^2} &\stackrel{L'H}{=} \lim_{x \rightarrow 0^+} \frac{\cot(x) - 1/x}{2x} = \lim_{x \rightarrow 0^+} \frac{x \cot(x) - 1}{2x^2} \\ &\stackrel{L'H}{=} \lim_{x \rightarrow 0^+} \frac{\cot(x) - x \csc^2(x)}{4x} \stackrel{L'H}{=} \lim_{x \rightarrow 0^+} \frac{1}{4} (-\csc^2(x) - \csc^2(x) + 2x \cot(x) \csc^2(x)) \\ &= \lim_{x \rightarrow 0^+} \frac{1}{4} \left(2x \frac{\cos(x)}{\sin^3(x)} - 2 \right) \frac{1}{\sin^3(x)} = \lim_{x \rightarrow 0^+} \frac{2x \cos(x) - 2 \sin(x)}{4 \sin^3(x)} \stackrel{L'H}{=} \lim_{x \rightarrow 0^+} \frac{-2x \sin(x)}{12 \sin^2(x) \cos(x)} \\ &= -\frac{1}{6} \lim_{x \rightarrow 0^+} \frac{x}{\sin(x) \cos(x)} \stackrel{L'H}{=} -\frac{1}{6} \lim_{x \rightarrow 0^+} \frac{1}{\cos^2(x) - \sin^2(x)} = -1/6. \text{ As } e^x \text{ is continuous,} \\ \lim_{x \rightarrow 0^+} \left(\frac{\sin(x)}{x} \right)^{1/x^2} &= \frac{1}{e^{1/6}} \end{aligned}$$

Q4 The value of k such that the equation $e^{2x} = k\sqrt{x}$ has one solution is

$$k = 2\sqrt{e}$$



Key idea: One solution means derivatives are also equal!

$$\begin{aligned} e^{2x} &= k\sqrt{x} \\ 2e^{2x} &= \frac{k}{2\sqrt{x}} \Rightarrow e^{2x} = \frac{k}{4\sqrt{x}} \Rightarrow x = 1/4 \\ e^{1/2} &= k \frac{1}{2} \Rightarrow k = 2\sqrt{e} \end{aligned}$$

Part 2. (10 points) Consider the two power series given by

$$J_0(x) = \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n}}{2^{2n} (n!)^2} \quad \text{and} \quad J_1(x) = \sum_{n=1}^{\infty} \frac{(-1)^n x^{2n+1}}{n! (n+1)! 2^{2n+1}}$$

These are called Bessel Functions of the first kind.

Q1 Determine the radii of convergence of $J_0(x)$ and $J_1(x)$.

i) $J_0(x) = \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n}}{(n!)^2 4^n}$

Then, $\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \lim_{n \rightarrow \infty} \left| \frac{(n!)^2 4^n}{(n+1)^2 (n!)^2 4^{n+1}} x^2 \right| = \lim_{n \rightarrow \infty} \frac{1}{4(n+1)^2} x^2 = 0 \quad \forall x \in \mathbb{R}.$

ROC = ∞ .

ii) Same idea for $J_1(x)$ to see ROC = ∞

Q2 Show that $y(x) = J_0(x)$ solves the differential equation

$$x^2 y''(x) + xy'(x) + x^2 y(x) = 0.$$

$J_0(x) = \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n}}{(n!)^2 4^n} = y(x)$ Set $a_n = \frac{(-1)^n}{(n!)^2 4^n}$

i) $x^2 y(x) = \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n+2}}{(n!)^2 4^n}$

ii) $xy'(x) = x \sum_{n=0}^{\infty} \frac{(-1)^n (2n) x^{2n-1}}{(n!)^2 4^n} = \sum_{n=0}^{\infty} \frac{(-1)^n (2n) x^{2n}}{(n!)^2 4^n}$

iii) $x^2 y''(x) = x^2 \sum_{n=0}^{\infty} \frac{(-1)^n (2n)(2n-1) x^{2n-2}}{(n!)^2 4^n} = \sum_{n=0}^{\infty} \frac{(-1)^n (2n)(2n-1) x^{2n}}{(n!)^2 4^n}$

Summing (ii) and (iii) = $\sum_{n=0}^{\infty} 4n^2 a_n x^{2n}$

Summing the above result with (i) yields

$$\underbrace{\sum_{n=0}^{\infty} a_n x^{2n+2}}_{= \sum_{n=1}^{\infty} a_{n-1} x^{2n}} + \sum_{n=0}^{\infty} 4n^2 a_n x^{2n}$$


hence consider $4n^2 a_n + a_{n-1}$

$$= \frac{4n^2 (-1)^n}{(n!)^2 4^n} + \frac{(-1)^{n-1}}{((n-1)!)^2 4^{n-1}} = 0.$$

Part 3 (15 points) Evaluate the following integrals.

Q1 $\int_0^1 \tan(\sin^{-1}(x)) dx =$

1



$$\tan(\sin^{-1}(x)) = \frac{x}{\sqrt{1-x^2}}$$

$$\int \tan(\sin^{-1}(x)) dx = \int \frac{x dx}{\sqrt{1-x^2}}$$

$$u = 1-x^2$$

$$du = -2x dx$$

$$= -\frac{1}{2} \int u^{-1/2} du = -u^{1/2} = -\sqrt{1-x^2}$$

$$\int_0^1 \tan(\sin^{-1}(x)) dx = -\sqrt{1-x^2} \Big|_0^1 = -(0-1) = 1$$

Q2 $\int \sqrt{1-x^2} dx =$

$$\frac{1}{2} x \sqrt{1-x^2} + \frac{1}{2} \arcsin(x)$$

$$\int \sqrt{1-x^2} dx$$

$$u = \sqrt{1-x^2} \quad dv = dx$$

$$du = \frac{-x}{\sqrt{1-x^2}} dx \quad v = x$$

$$= x \sqrt{1-x^2} + \int \frac{x^2}{\sqrt{1-x^2}} dx$$

$$\int \frac{x^2}{\sqrt{1-x^2}} dx = \int \frac{x^2-1+1}{\sqrt{1-x^2}} dx$$

$$= -\int \frac{1-x^2}{\sqrt{1-x^2}} dx + \int \frac{dx}{\sqrt{1-x^2}}$$

$$= -\int \sqrt{1-x^2} dx + \arcsin(x)$$

$$\Rightarrow \underbrace{\int \sqrt{1-x^2} dx = -\int \sqrt{1-x^2} dx + x \sqrt{1-x^2} + \arcsin(x)}_{\text{combine to } 2 \int \sqrt{1-x^2} dx} \Rightarrow \int \sqrt{1-x^2} dx = \frac{1}{2} x \sqrt{1-x^2} + \frac{1}{2} \arcsin(x)$$

Q3 $\int \frac{\cot(x)}{\ln(\sin(x))} dx =$

$$\ln(\ln(\sin(x)))$$

$$\int \frac{\cot(x)}{\ln(\sin(x))} dx = \int \frac{\cos(x)}{\sin(x) \ln(\sin(x))} dx$$

$$u = \sin(x)$$

$$du = \cos(x) dx$$

$$= \int \frac{du}{u \ln(u)}$$

$$s = \ln(u)$$

$$ds = \frac{du}{u}$$

$$= \int \frac{ds}{s} = \ln(s) = \ln(\ln(\sin(x)))$$

Part 4. (15 points)

Q1 If $f(x) = \sum_{n=1}^{\infty} a_n x^n$, then $\sum_{n=1}^{\infty} n^2 a_n x^n =$

$$x f'(x) + x^2 f''(x)$$

$$f(x) = \sum_{n=1}^{\infty} a_n x^n, \quad f'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1} \Rightarrow x f'(x) = \sum_{n=1}^{\infty} n a_n x^n$$

$$\Rightarrow \frac{d}{dx} (x f'(x)) = f'(x) + x f''(x) = \sum_{n=1}^{\infty} n^2 a_n x^{n-1}$$

$$\Rightarrow x f'(x) + x^2 f''(x) = \sum_{n=1}^{\infty} n^2 a_n x^n$$

Q2 The values of p and q such that $\sum_{n=3}^{\infty} \frac{1}{n^p \ln^q(n)}$ converges are

$$p=1, q>1$$

$$p>1, q \in \mathbb{R}$$

$$\frac{p=1}{\int_3^{\infty} \frac{dx}{x \ln^q(x)}} = \int_3^{\infty} \frac{dx}{u^q}$$

converges if and
and only if $q>1$.

$$p=1, q>1$$

$$\frac{p>1}{n^p \ln^q(n) > n^p} \Rightarrow \frac{1}{n^p \ln^q(n)} < \frac{1}{n^p}$$

converges
via p-series

Direct comparison test
yields convergence for

$$p>1, q \in \mathbb{R}$$

$$\frac{p<1}{\sum_{n=3}^{\infty} \frac{1}{n^p \ln^q(n)}}$$

$$\int_3^{\infty} \frac{dx}{x^p \ln^q(x)}$$

$$u = \ln(x) \Rightarrow x = e^u$$

$$du = \frac{dx}{x}$$

$$= \int_{\ln(3)}^{\infty} \frac{e^{(1-p)u}}{u^q} du \rightarrow +\infty \quad \forall p \in (0,1), q \in \mathbb{R}$$

Q3 Evaluate the following integral: $\int \frac{\ln(\ln(x))}{x} dx =$

$$\ln(x) \ln(\ln(x)) - \ln(x)$$

$$\int \frac{\ln(\ln(x))}{x} dx$$

$$t = \ln(x)$$

$$dt = \frac{dx}{x}$$

$$= \int \ln(t) dt$$

$$u = \ln(t) \quad dv = dt$$

$$du = \frac{dt}{t} \quad v = t$$

$$= t \ln(t) - \int dt$$

$$= t \ln(t) - t$$

$$= \ln(x) \ln(\ln(x)) - \ln(x)$$

Part 5. (15 points)**Q1** Consider the parametric equations

$$x(t) = e^t \cos(t) \quad y(t) = e^t \sin(t)$$

Eliminate the parameter and find $\frac{dy}{dx}$ in terms of t .

$$\begin{aligned} \text{i)} \quad x^2 + y^2 &= e^{2t} \cos^2(t) + e^{2t} \sin^2(t) \\ &= e^{2t} \\ \Rightarrow t &= \frac{1}{2} \ln(x^2 + y^2) \\ \text{ii)} \quad \frac{x}{y} &= \frac{e^t \cos(t)}{e^t \sin(t)} = \cot(t) \\ \Rightarrow t &= \cot^{-1}(x/y) \\ \cot^{-1}(x/y) &= \frac{1}{2} \ln(x^2 + y^2) \end{aligned}$$

$$\frac{dy}{dx} = \frac{dy/dt}{dx/dt} = \frac{e^t \sin(t) + e^t \cos(t)}{e^t \cos(t) - e^t \sin(t)} = \frac{y+x}{x-y}$$

Q2 Find the length of the polar curve of $r = \frac{1}{\theta}$ from $\theta = \frac{\pi}{2}$ to $\theta = 2\pi$

$$\begin{aligned} r(\theta) &= \frac{1}{\theta} \Rightarrow r^2 + (r'(\theta))^2 = \frac{1}{\theta^2} + \frac{1}{\theta^4} \\ r'(\theta) &= \frac{-1}{\theta^2} \quad \sqrt{\frac{1}{\theta^2} + \frac{1}{\theta^4}} = \frac{\sqrt{\theta^2 + 1}}{\theta^2} \\ \int \sqrt{r^2 + (r'(\theta))^2} d\theta &= \int \frac{\sqrt{\theta^2 + 1}}{\theta^2} d\theta = -\frac{\sqrt{1+\theta^2}}{\theta} + \int \frac{d\theta}{\sqrt{1+\theta^2}} \\ u &= \sqrt{1+\theta^2} \quad dv = \frac{d\theta}{\theta^2} \\ du &= \frac{\theta d\theta}{\sqrt{1+\theta^2}} \quad v = -\theta^{-1} \\ &= \ln(\theta + \sqrt{1+\theta^2}) - \frac{\sqrt{1+\theta^2}}{\theta} \end{aligned}$$

Substituting yields

$$\begin{aligned} &= \int \frac{dt}{1-t^2} = \int \frac{A}{1-t} dt + \int \frac{B}{1+t} dt \\ &= \frac{1}{2} \int \frac{dt}{1-t} + \frac{1}{2} \int \frac{dt}{1+t} = \frac{1}{2} \ln \left| \frac{1+t}{1-t} \right| \\ &= \frac{1+t}{1-t} \cdot \frac{(1+t)}{(1+t)} = \frac{(1+t)^2}{1-t^2} = -\theta^2(1+t^2) \\ &= \ln(\theta(1+t)) = \ln(\theta + \theta\sqrt{1+\theta^2}) \\ &= \ln(\theta + \sqrt{1+\theta^2}) \end{aligned}$$

Q3 Eliminate the parameter of the parametric equations $x = 2 \cos(\theta)$, $y = 1 + \sin(\theta)$. Find the points of the horizontal and vertical tangent lines.

$$\begin{aligned} x &= 2 \cos(\theta) \Rightarrow \cos(\theta) = x/2 \\ y &= 1 + \sin(\theta) \Rightarrow \sin(\theta) = y-1 \\ \text{Hence, } x^2/4 + y^2 - 2y + 1 &= 1. \end{aligned}$$

$$\begin{aligned} \frac{dx}{d\theta} &= -2 \sin(\theta) \\ -2 \sin(\theta) &= 0 \\ \theta &= 0, \pi \end{aligned}$$

$$\begin{aligned} \frac{dy}{d\theta} &= \cos(\theta) \\ \cos(\theta) &= 0 \\ \theta &= \pi/2, 3\pi/2 \end{aligned}$$

Part 6. (10 points)

Q1 The general solution to $\frac{dy}{dx} = \frac{3^x y (\ln^2(y) + 3 \ln(y) + 2)}{1 + 9^x}$ is

$$\frac{\ln(y)+1}{\ln(y)+2} = e^{\tan^{-1}(3^x)} + C$$

$$\frac{dy}{dx} = \frac{3^x y (\ln^2(y) + 3 \ln(y) + 2)}{1 + 9^x}$$

$$\int \frac{dy}{y(\ln^2(y) + 3 \ln(y) + 2)} = \int \frac{3^x}{1 + (3^x)^2} dx$$

$$u_1 = \ln(y) \quad u_2 = 3^x$$

$$du_1 = \frac{1}{y} dy \quad du = \ln(3) 3^x dx$$

$$\int \frac{du_1}{(u_1+1)(u_1+2)} = \frac{1}{\ln(3)} \int \frac{du_2}{1+(u_2)^2}$$

$$\int \frac{-1}{u_1+2} + \frac{1}{u_1+1} du = \frac{1}{\ln(3)} \int \frac{du_2}{1+(u_2)^2}$$

$$-\ln(u_1+2) + \ln(u_1+1) = \frac{1}{\ln(3)} \tan^{-1}(u_2) + C$$

$$-\ln(\ln(y)+2) + \ln(\ln(y)+1) = \frac{1}{\ln(3)} \tan^{-1}(3^x) + C$$

$$\frac{1}{(u_1+2)(u_1+1)} = \frac{A}{u_1+2} + \frac{B}{u_1+1}$$

$$A = -1, B = 1$$

Q2 Use power series to evaluate: $\lim_{x \rightarrow 0} \frac{1 - \cos(x)}{1 + x - e^x} =$

-1

$$\begin{aligned} \lim_{x \rightarrow 0} \frac{1 - \cos(x)}{1 + x - e^x} &= \lim_{x \rightarrow 0} \frac{1 - \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n}}{(2n)!}}{1 + x + \sum_{n=0}^{\infty} \frac{x^n}{n!}} = \lim_{x \rightarrow 0} \frac{1 - (1 - \frac{x^2}{2} + \frac{x^4}{4!} - \dots)}{1 + x - (1 + x + \frac{x^2}{2} + \dots)} \\ &= \lim_{x \rightarrow 0} \frac{\frac{x^2}{2} - \frac{x^4}{4!} + \dots}{-x^2/2 - x^3/3! - x^4/4! - \dots} \stackrel{L'H}{=} \lim_{x \rightarrow 0} \frac{x - x^3/3! + x^5/5! - \dots}{-x - x^2/2! - x^3/3! - \dots} \stackrel{L'H}{=} \lim_{x \rightarrow 0} \frac{1 - x^2/2! + x^4/4! - \dots}{-1 - x - x^2/2! - \dots} \\ &= -1 \end{aligned}$$

Part 7 (20 points) Determine if the following series converge or diverge. Show your work and state any necessary hypotheses.

$$\text{Q1 } \sum_{n=0}^{\infty} \frac{1}{n\sqrt{n^2+1}}$$

Method 1 (Direct Comparison Test)
 $n^2+1 > n^2$ As $\frac{1}{n^2}$ converges via the
 $\Rightarrow \sqrt{n^2+1} > n$ p-series test, the direct
 $\Rightarrow n\sqrt{n^2+1} > n^2$ comparison test yields
 $\Rightarrow \frac{1}{n\sqrt{n^2+1}} < \frac{1}{n^2}$ $\frac{1}{n\sqrt{n^2+1}}$ converges, too.

Method 2 (Limit Comparison Test)
 Set $a_n = \frac{1}{n\sqrt{n^2+1}}$ and $b_n = \frac{1}{n^2}$. As
 $\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = 1 > 0$, the limit comparison test
 yields that $\sum a_n$ converges as $\sum b_n$
 converges.

$$\text{Q2 } \sum_{n=1}^{\infty} \frac{(n!)^n}{n^{4n}}$$

$$\begin{aligned} \text{Set } a_n &= \frac{(n!)^n}{n^{4n}}. \text{ Then, } \left| \frac{a_{n+1}}{a_n} \right| = \left| \frac{(n+1)^n (n!)^n}{(n+1)^{4n+4}} \cdot \frac{n^{4n}}{(n!)^n} \right| \\ &= \left| \frac{n^{4n}}{(n+1)^{3n+4}} \right| = \left| \frac{n^n \cdot n^{3n}}{(n+1)^4 (n+1)^{3n}} \right| = \frac{n^n}{(n+1)^4} \left(\frac{n}{n+1} \right)^{3n} \\ &\quad \rightarrow \infty \quad \rightarrow 1/e^3 \end{aligned}$$

By the ratio test yields divergence

$$\text{Q3 } \sum_{n=1}^{\infty} (-1)^n \frac{n^2}{2^n} \text{ (Use the alternating series test)}$$

$$\begin{aligned} \text{Set } a_n &= \frac{n^2}{2^n} \text{ and consider } f(x) = \frac{x^2}{2^x}. \text{ Then, } \lim_{x \rightarrow \infty} f(x) = \lim_{x \rightarrow \infty} \frac{x^2}{2^x} \\ &\stackrel{\text{L'H}}{=} \lim_{x \rightarrow \infty} \frac{2x}{\ln(2)2^x} \stackrel{\text{L'H}}{=} \lim_{x \rightarrow \infty} \frac{2}{\ln^2(2)2^x} = 0. \text{ Hence, } \lim_{n \rightarrow \infty} a_n = 0. \text{ Thus, the Alternating} \end{aligned}$$

Series Test yields convergence.

$$\text{Q4 } \sum_{n=1}^{\infty} \sin\left(\frac{n\pi}{6}\right)$$

$\lim_{n \rightarrow \infty} \sin\left(\frac{n\pi}{6}\right) = \text{DNE}$. By the
 divergence test $\sum \sin\left(\frac{n\pi}{6}\right)$ diverges.

Part 8 (20 points) Define the function $f(x) = \int_0^x \frac{1}{(t+1)\sqrt{1-\ln^2(t+1)}} dt$

Q1 The domain of $f(x)$ is

$$\{x \in \mathbb{R} \mid e^{-1}-1 \leq x \leq e-1\}$$

$$f(x) = \int_0^x \frac{dt}{(t+1)\sqrt{1-\ln^2(t+1)}} \stackrel{u=t+1}{=} \int_0^x \frac{du}{u\sqrt{1-\ln^2(u)}} \stackrel{s=\ln(u)}{=} \int_0^x \frac{ds}{\sqrt{1-s^2}} = \sin^{-1}(\ln(x+1))$$

$$f(x) = \arcsin(\ln(x+1))$$

$$-1 \leq \ln(x+1) \leq 1$$

$$e^{-1}-1 \leq x \leq e-1$$

Q2 $f(x)$ has an inverse on

$$\{x \in \mathbb{R} \mid e^{-1}-1 \leq x \leq e-1\}$$

$$f'(x) = \frac{1}{(x+1)\sqrt{1-\ln^2(x+1)}} = \underbrace{\frac{1}{x+1}}_{< 0} \cdot \underbrace{\frac{1}{\sqrt{1-\ln^2(x+1)}}}_{< 0}$$

Q3 $f\left(\sum_{n=1}^{\infty} \frac{(3)^{n/2}}{n! 2^n}\right) = \frac{\pi}{3}$

$$\sum_{n=0}^{\infty} \frac{3^{n/2}}{n! 2^n} = \sum_{n=0}^{\infty} \frac{(\sqrt{3}/2)^n}{n!} = e^{\sqrt{3}/2}$$

$$\Rightarrow \sum_{n=1}^{\infty} \frac{3^{n/2}}{n! 2^n} = e^{\sqrt{3}/2} - 1$$

$$\begin{aligned} f(e^{\sqrt{3}/2} - 1) &= \arcsin(\ln(e^{\sqrt{3}/2} - 1 + 1)) \\ &= \sin^{-1}(\sqrt{3}/2) = \pi/3 \end{aligned}$$

Q4 $(f^{-1})'(1/2) =$

$$e^{\sin(1/2)} \cos(1/2)$$

$$(f^{-1})'(\frac{1}{2}) = \frac{1}{f'(f^{-1}(1/2))} = \frac{1}{e^{\sin(1/2)} \sqrt{1-\sin^2(1/2)}} = e^{\sin(1/2)} \cos(1/2)$$

$$\arcsin(\ln(x+1)) = 1/2$$

$$\ln(x+1) = \sin(1/2)$$

$$x = e^{\sin(1/2)} - 1$$

Part 9 (20 points) Suppose that $y = f(x)$ is a solution to the differential equation

$$\frac{dy}{dx} = yx \ln(x) + k \sin(x)$$

where $k > 0$ and $f(1) = 4$.

Q1 The second order Taylor polynomial of $f(x)$ at $x = 1$ is given by

$$f(x) \approx 4 + k \sin(1)(x-1) + \frac{1}{2}(4 + k \cos(1))(x-1)^2$$

$$f(1) = 4$$

$$f'(x) = \frac{dy}{dx} = yx \ln(x) + k \sin(x)$$

$$f'(1) = (4)(1) \ln(1) + k \sin(1) = k \sin(1)$$

$$f''(x) = \frac{dy}{dx} x \ln(x) + y (\ln(x) + x) + k \cos(x)$$

$$f''(1) = (k \sin(1))(1) \ln(1) + 4(\ln(1) + 1) + k \cos(1) = 4 + k \cos(1)$$

$$f(x) \approx 4 + k \sin(1)(x-1) + \frac{1}{2}(4 + k \cos(1))(x-1)^2$$

Q2 If $k = 0$, then the solution to the initial value problem is given by

$$y = \exp\left(\frac{x^2}{2}(\ln(x) - \frac{1}{2}) + \ln(4) + \frac{1}{4}\right)$$

$$\text{If } k=0, \quad \frac{dy}{dx} = yx \ln(x) \quad c = \ln(4) + \frac{1}{4}$$

$$\frac{dy}{y} = x \ln(x) dx$$

$$\ln|y| = \frac{x^2}{2}(\ln(x) - \frac{1}{2}) + \ln(4) + \frac{1}{4}$$

$$\int \frac{dy}{y} = \int x \ln(x) dx$$

$$y = \exp\left(\frac{x^2}{2}(\ln(x) - \frac{1}{2}) + \ln(4) + \frac{1}{4}\right)$$

$$u = \ln(x) \quad dv = x dx$$

$$du = \frac{dx}{x} \quad v = \frac{x^2}{2}$$

$$\ln|y| = \frac{x^2}{2} \ln(x) - \frac{1}{2} \int x dx$$

$$\ln|y| = \frac{x^2}{2}(\ln(x) - \frac{1}{2}) + C$$

$$f(1) = 4$$

$$\ln(4) = \frac{1}{4}(-1) + C$$

Q3 The domain of $f(x)$ is

$$x > 0$$

Part 10 (25 points) Define the function $g(x) = xe^x$.

Q1 Calculate $\int_0^1 g(x) dx$ in two ways: an integration technique and a power series technique.

$$\begin{aligned} \text{i) } \int_0^1 x e^x dx \\ u = x \quad dv = e^x dx \\ du = dx \quad v = e^x \\ = x e^x \Big|_0^1 - \int_0^1 e^x dx \\ = e - (e - 1) = 1 \end{aligned}$$

$$\begin{aligned} \text{ii) } x e^x &= x \sum_{n=0}^{\infty} \frac{x^n}{n!} = \sum_{n=0}^{\infty} \frac{x^{n+1}}{n!} \\ \int_0^1 x e^x dx &= \int_0^1 \sum_{n=0}^{\infty} \frac{x^{n+1}}{n!} dx = \sum_{n=0}^{\infty} \int_0^1 \frac{x^{n+1}}{n!} dx \\ &= \sum_{n=0}^{\infty} \frac{1}{(n+2)n!} \end{aligned}$$

Q2 $y = g(x)$ satisfies the following differential equation

$$\frac{dy}{dx} = x e^x + e^x$$

$$\frac{dy}{dx} = y + \frac{y}{x} = \left(\frac{x+1}{x} \right) y$$

$$\frac{dy}{dx} = \left(\frac{x+1}{x} \right) y$$

Q3 The above differential equation is **SEPARABLE** or **NON-SEPARABLE**.

Q4 The domain of $h(p) = \int_1^{\infty} \frac{g(x)}{e^{px}} dx$ is

$$p > 1$$

$$h(p) = \int_1^{\infty} \frac{g(x)}{e^{px}} dx = \int_1^{\infty} x e^{(1-p)x} dx = \lim_{c \rightarrow \infty} \underbrace{\frac{1}{p-1} x e^{(1-p)x} \Big|_1^c - \frac{1}{1-p} \int_1^c e^{(1-p)x} dx}_{\text{converges if and only if } p > 1}$$

$$\begin{aligned} u = x \quad dv = e^{(1-p)x} dx \\ du = dx \quad v = \frac{1}{1-p} e^{(1-p)x} \end{aligned}$$

converges if and only if $p > 1$.

Q5 $g(x)$ has an inverse on the domain

$$(-\infty, -1) \cup (-1, \infty)$$

$$g'(x) = (x+1)e^x$$

$$g'(x) > 0 \quad \forall x > -1$$

$$g'(x) < 0 \quad \forall x < -1$$


As $e^x > 0 \quad \forall x \in \mathbb{R}$, the sign of $x+1$ determines if $g(x)$ has an inverse.

Q6 $\int_0^e g^{-1}(x) dx =$ $e-1$. (Hint: what is an inverse?)

$$\text{triangle area} = \frac{1}{2} e^2$$

$$\begin{aligned} \text{area of orange region} &= \int_0^1 g(x) - x dx + \int_1^e e - x dx \\ &= 1 - \frac{1}{2} + e(e-1) - \frac{1}{2}(e^2-1) \\ &= 1 + \frac{1}{2} e^2 - e \end{aligned}$$

$$\Rightarrow \int_0^e g^{-1}(x) dx = \frac{1}{2} e^2 - (1 + \frac{1}{2} e^2 - e) = e - 1$$

Part 11 *Extra space for work*

End of Exam. Check your work!